



Achilles Tendon Repair Rehabilitation

PHASE 1 (Immediate Post-Operative Phase): Generally 0-6 Weeks Post-Op

Rehabilitation Goals

- Protect the surgical repair
- Avoid a “stiff” ankle
- Attain DF ROM to neutral at 6 weeks post-op
- Minimize pain, swelling, muscle atrophy, and deconditioning
- Independent gait without assistive device

PRECAUTIONS

- **ALWAYS** wear ankle CAM boot for ambulation
- Limit ankle DF to 0° until 6 weeks post-op

Crutches/Brace

Per Ortho Surgeon* (typically):

- Weeks 1-2: 1/2-inch heel lift; toe touch WB @ 0-10% body weight while in post-op splint for 10 days; progress to WBAT
- Weeks 3-4: 1/4-inch heel lift; WBAT
- Weeks 5-6: 1/8-inch heel lift as needed; WBAT ***NOTE: May progress earlier based on Ortho preference
- D/C crutches when gait is normalized; goal is between 4-6 weeks post-op

Wound

- Shower after post-op day 2 (cover splint/cast when showering)
- DO NOT SUBMERGE ankle in tub or pool for 4 weeks
- Suture removal @ 10-14 days post-op per Ortho
- Begin scar massage after incision site has healed and scar is formed

Rehabilitation

- Keep LE elevated as much as possible; ice ankle when applicable
- Ankle pumps while in splint
- Begin exercises listed below



Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center

410-539-2227 410-539-2240

301 St. Paul Pl. Baltimore MD 21202

Achilles Tendon Repair Rehabilitation

PHASE 1 (Immediate Post-Operative Phase): Generally 0-6 Weeks Post-Op

Weeks 1-2

- Hip and knee AROM exercises
- Quad sets and glute sets
- Intrinsic foot strengthening/toe posture and short foot exercises (i.e. resisted towel curls and toe yoga)
 - Knee and hip supine/seated open kinetic chain (OKC) strengthening exercises as tolerated (i.e. SLRs, LAQs, and SAQs)
- Ankle isometrics as tolerated
- Non-resisted active calf pumps (from neutral to PF as tolerated) once splint is removed (i.e. 50-100 reps, 5-6x per day)
- HS stretching

Weeks 3-4

- UBE for aerobic strength/endurance and seated UE weight lifting
- Recumbent bike as tolerable while donning CAM boot
- Intrinsic foot strengthening/toe posture and short foot exercises
- Gentle, seated Achilles towel stretch (pain-free at tendon)
- Ankle ROM exercises (i.e. ankle pumps, alphabets, and CW/CCW circles)
- Ankle isometrics as tolerated
- Non-resisted active calf pumps (from neutral to PF as tolerated) once splint is removed (i.e. 50-100 reps, 5-6x per day)
- Knee and hip supine/seated OKC strengthening exercises as tolerated (i.e. resisted knee extensions, HS curls, and hip strengthening)
- LE stretching: HS, glutes, ITB, piriformis, and quads
- AAROM self-mobs for PF
- OKC proprioceptive exercises
- Manual Therapy: ankle/foot mobilizations as needed; limit DF to 0° until 6 weeks post-op

Weeks 5-6

- Low intensity stationary bike with light resistance (5-10 minutes)
- Pain-free ankle isometrics
- Active calf pumps (from neutral to PF as tolerated) once splint is removed (i.e. 50-100 reps, 5-6x per day)
- AAROM self-mobs for PF
- Ankle strengthening with light tubing all directions as tolerated
- Seated SL heel raises
- Continue hip and knee supine/seated OKC strengthening exercises as tolerable (i.e. resisted knee extensions, HS curls, and hip strengthening)
- Manual Therapy: ankle/foot mobilizations as needed; limit DF to 0° until 6 weeks post-op
- Beginner-level pool exercises (after incision is healed)
- Chest-deep water walking and exercises (within precautions)

Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center



410-539-2227



410-539-2240

301 St. Paul Pl. Baltimore MD 21202

Achilles Tendon Repair Rehabilitation

PHASE 1 (Immediate Post-Operative Phase): Generally 0-6 Weeks Post-Op

Follow Up

- Supervised rehab: 1-2x per week
- PT re-eval: every 2-4 weeks as needed
- Ortho re-eval: 2 weeks post-op and 6 weeks post-op

Documentation

- Precautions, pain level, medications, and modalities
- Ankle ROM & gait
- Observation
 - Incision sites healing well?
 - Signs/symptoms of infection?
 - Effusion? - Neurovascular status
 - Distal pulses?
 - Motor/sensation nerves intact?
 - Presence of calf pain?

Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center

410-539-2227 410-539-2240

301 St. Paul Pl. Baltimore MD 21202



Achilles Tendon Repair Rehabilitation

PHASE 2: Generally 7-12 Weeks Post-Op

Rehabilitation Goals

- Full ankle ROM
- Independent ambulation with level walking and stair negotiation without any observed gait deviation

PRECAUTIONS

- **NO RUNNING** until after 4 months post-op

Cryotherapy

- Cold with compression/elevation (i.e. ice with compression wrap)

Rehabilitation

- Continue Phase 1 exercises and scar massage as needed
- Progress to the following exercises and increase intensity gradually as tolerated (i.e. minimal to no increase in ankle pain, stiffness, or edema since the previous exercise session)
- All strengthening should be done starting with low weights and high reps before progressing resistance



Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center

410-539-2227 410-539-2240

301 St. Paul Pl. Baltimore MD 21202

Achilles Tendon Repair Rehabilitation

PHASE 2: Generally 7-12 Weeks Post-Op

Weeks 7-8

- Stationary bike for conditioning; resistance as tolerated
- Ankle ROM exercises (add mobilizations/manual stretching as needed)
- Seated wobble board
- Gait training (i.e. cone taps, marching, retro-walking, cariocas, and shuffles)
- Ankle strengthening with tubing - DL squats to depth of tolerance (add resistance gradually)
- Forward, lateral, and retro step-ups (start with 4" step and progress as tolerated)
- Standing DL heel raises
- Standing gastroc/soleus stretches
- Progressive hip/knee strengthening (i.e. knee extensions, leg press, HS curls, and hip ABD/ADD) -
Manual Therapy: ankle mobilizations in all directions until ROM is WNL

Weeks 9-12

- UBE for aerobic strength/endurance and seated UE weight lifting
- Recumbent bike as tolerable while donning CAM boot
- Intrinsic foot strengthening/toe posture and short foot exercises
- Gentle, seated Achilles towel stretch (pain-free at tendon)
- Ankle ROM exercises (i.e. ankle pumps, alphabets, and CW/CCW circles)
- Ankle isometrics as tolerated
- Non-resisted active calf pumps (from neutral to PF as tolerated) once splint is removed (i.e. 50-100 reps, 5-6x per day)
- Knee and hip supine/seated OKC strengthening exercises as tolerated (i.e. resisted knee extensions, HS curls, and hip strengthening)
- LE stretching: HS, glutes, ITB, piriformis, and quads
- AAROM self-mobs for PF
- OKC proprioceptive exercises
- Manual Therapy: ankle/foot mobilizations as needed; limit DF to 0° until 6 weeks post-op

Weeks 5-6

- Elliptical/stationary bike with progressive resistance
- Progress DL squats as tolerated; add resistance gradually
- Progress forward, lateral, and retro step-ups as tolerated
- Standing heel raise progression from DL → SL as able
- Progressive hip/knee strengthening (i.e. knee extensions, leg press, HS curls, and hip ABD/ADD) -
Standing balance progression from DL → SL as able
- Standing gastroc/soleus stretches - Intermediate-level pool exercises

Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center

410-539-2227 410-539-2240

301 St. Paul Pl. Baltimore MD 21202

Achilles Tendon Repair Rehabilitation

PHASE 2: Generally 7-12 Weeks Post-Op

Follow Up

- Supervised rehab: 1-3x per week as needed
- PT re-eval: every 2-4 weeks as needed
- Ortho re-eval: 3 months post-op

Documentation

- Precautions, pain level, medications, and modalities
- Ankle ROM, strength, and gait
- Observation
 - Incision sites healing well?
 - Signs/symptoms of infection?
 - Effusion?

Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center

410-539-2227 410-539-2240

301 St. Paul Pl. Baltimore MD 21202



Achilles Tendon Repair Rehabilitation

PHASE 3: Generally 4-6 Months Post-Op

Rehabilitation Goals

- Full ankle ROM
- Perform 20+ SL heel raises to $\geq 75\%$ height of contralateral limb
- Hop for distance $> 90\%$ of uninvolved side
- Jog at own pace/distance without pain
- Meet occupational requirements at 6 months

PRECAUTIONS

- **NO RUNNING** until after 4 months post-op

Crutches/Brace

- Ankle lace-up brace as needed

Rehabilitation

- Continue Phase 2 exercises as needed
- Progress to the following exercises, gradually increasing intensity and duration as tolerated, as long as there is no increased ankle pain or edema



Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center

410-539-2227 410-539-2240

301 St. Paul Pl. Baltimore MD 21202

Achilles Tendon Repair Rehabilitation

PHASE 3: Generally 4-6 Months Post-Op

Weeks 13-16

- Gradually add stairmaster to low-impact conditioning regimen
- Progressive LE strengthening (i.e. calf press, leg press, squats, HS curls, and hip ABD/ADD)
- Light agility exercises as tolerated (i.e. fitter, slide board, figure 8s, gentle loops, large zig zags, and agility ladder)
- Progressive balance training as needed
- Progressive pool program as tolerated

Weeks 17-24

- Progressive jogging program
- May begin each session with jogging on treadmill for 5 minutes
- Increase time and/or distance no more than 10-20% per week
- Progressive agility/functional training
- Begin at 25-50% intensity and progress gradually
- Jumping, hopping, directional jogging, cariocas, shuffles, etc.

Follow Up

- Supervised rehab: 1-2x per week as needed
- PT re-eval: monthly
- Ortho re-eval: 6 months post-op

Documentation

- Pain level and medications
- Ankle ROM and strength
- Hop test for distance
- Functional activity tolerance (i.e. stairs, jogging)

Discharge Goals

- Hop test and Y-balance limb symmetry > 90%
- Isokinetic testing limb symmetry > 85%
- Hand held dynamometer > 85% for PF
- Injury-Psychological Readiness to Return to Sport Scale (> 50% indicates readiness)

Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center

410-539-2227 410-539-2240

301 St. Paul Pl. Baltimore MD 21202



Achilles Tendon Repair Rehabilitation

PHASE 3: Generally 4-6 Months Post-Op

Miscellaneous

- After 6 months post-op, exercises in Phase 3 are continued and gradually increasing in intensity and duration as tolerated
- Pass Service fitness test at 6-9 months
- Progress activities for return to sport/collision sports or aggressive military training (i.e. airborne school) based on the patient's functional performance and endurance. This time period will be directed by the Ortho Surgeon and the Physical Therapist. This may require between 9-12 months before cleared without restrictions.

Dr. Mark Slabaugh

Orthopedics and Joint Replacement | Mercy Medical Center

410-539-2227 410-539-2240

301 St. Paul Pl. Baltimore MD 21202